

**9th Class 2019**

Math (Science)	Group-I	Paper-I
Time: 20 Minutes	(Objective Type)	Max Marks: 15

Note: Four possible answers A, B, C and D to each question are given. The choice which you think is correct, fill that circle in front of that question with Marker or Pen ink in the answer-book. Cutting or filling two or more circles will result in zero mark in that question.

- 1-1- The square root of  $a^2 - 2a + 1$  is ----:
  - (a)  $\pm(a + 1)$
  - (b)  $\pm(a - 1) \checkmark$
  - (c)  $(a - 1)$
  - (d)  $(a + 1)$
- 2- A diagonal of a parallelogram divides it into --- congruent triangles:
  - (a) 2  $\checkmark$
  - (b) 3
  - (c) 4
  - (d) 6
- 3- Two parallel lines intersect at --- point / points:
  - (a) Three
  - (b) Two
  - (c) One
  - (d) No  $\checkmark$
- 4- The diagonals of parallelogram --- each other:
  - (a) Bisect  $\checkmark$
  - (b) Trisect
  - (c) Bisect at right angle
  - (d) Trisect at right angle
- 5- Order of transpose of matrix  $\begin{bmatrix} 2 & 1 \\ 0 & 1 \\ 3 & 2 \end{bmatrix}$  is:
  - (a) 3 - by - 2  $\checkmark$
  - (b) 2 - by - 3
  - (c) 1 - by - 3
  - (d) 3 - by - 1
- 6-  $x = 0$  is a solution of the inequality ----:
  - (a)  $x > 0$
  - (b)  $3x + 5 < 0$
  - (c)  $x + 2 < 0$
  - (d)  $x - 2 < 0 \checkmark$



- 7- Any point on the bisector of an angle is --- from its arms:
- (a) Un-equidistance
  - (b) Equidistance ✓
  - (c) Large distance
  - (d) Small distance
- 8-  $(3 + \sqrt{2})(3 - \sqrt{2})$  is equal to:
- (a)  $7\sqrt{\quad}$
  - (b)  $-7$
  - (c)  $-1$
  - (d)  $1$
- 9- Unit of ratio is ----:
- (a) Degree
  - (b) None ✓
  - (c) cm
  - (d)  $\pi$
- 10- Point  $(-3, -3)$  lies in quadrant:
- (a) I
  - (b) II
  - (c) III ✓
  - (d) IV
- 11- The logarithm of unity to any base is ---- :
- (a) 1
  - (b) 10
  - (c) e
  - (d)  $0\sqrt{\quad}$
- 12- A ray has end points:
- (a)  $1\sqrt{\quad}$
  - (b) 2
  - (c) 3
  - (d) 4
- 13- Distance between points  $(0, 0)$  and  $(1, 1)$  is ----:
- (a) 0
  - (b) 1
  - (c) 2
  - (d)  $\sqrt{2}\sqrt{\quad}$
- 14- The conjugate of  $5 + 4i$  is ---- :
- (a)  $-5 + 4i$
  - (b)  $-5 - 4i$
  - (c)  $5 - 4i\sqrt{\quad}$
  - (d)  $5 + 4i$
- 15- In a parallelogram opposite angles are ----:
- (a) Perpendicular
  - (b) Equal ✓
  - (c) Unequal
  - (d) Acute



## 9th Class 2019

Math (Science)	Group-I	Paper-I
Time: 2.10 Hours	(Subjective Type)	Max. Marks: 60

## (Part-I)

2. Write short answers to any Six (6) questions: 12

(i) Define column matrix.

**Ans** A matrix is called a column matrix if it has only one

column. e.g.,  $M = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $N = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$  are column matrices

of order 2 – by – 1 and 3 – by – 1 respectively.

(ii) Find the transpose of the matrix:  $B = [5 \ 1 \ -6]$

**Ans** Given,  $B = [5 \ 1 \ -6]$

Transpose:  $B^t = \begin{bmatrix} 5 \\ 1 \\ -6 \end{bmatrix}$

(iii) Simplify:  $\sqrt[3]{-125}$ .

**Ans**

$$\begin{aligned} \sqrt[3]{-125} &= (-125)^{1/3} \\ &= [-(5 \times 5 \times 5)]^{1/3} \\ &= [-(5^3)]^{1/3} \\ &= -5^{3 \times 1/3} = -5^1 \\ &= -5 \end{aligned}$$

(iv) Write real and imaginary parts of the number:

$$-1 + 2i$$

**Ans** Real Part = -1

Imaginary Part = 2i

(v) Express in scientific notation: 83,000

**Ans**  $83000 = 83000 \times \frac{10000}{10000}$

$$= \frac{83000}{10000} \times 10000$$

$$= 8.3 \times 10^4$$



(vi) Find the value of  $x$   $\log_3 x = 4$

**Ans**  $\log_3 x = 4$

By writing in exponential form, we have:

$$x = 3^4$$

Thus,  $x = 81$

(vii) Evaluate  $\frac{x^3y - 2z}{xz}$  for  $x = -1, y = -9, z = 4$

**Ans** By putting the values of  $x, y$  and  $z$  in the given expression, i.e.,

$$\begin{aligned} \frac{x^3y - 2z}{xz} &= \frac{(-1)^3(-9) - 2(4)}{(-1)(4)} \\ &= \frac{(-1)(-9) - 8}{-4} \\ &= \frac{9 - 8}{-4} \\ &= \frac{-1}{4} \end{aligned}$$

(viii) Rationalize the denominator:  $\frac{58}{7 - 2\sqrt{5}}$

**Ans**

$$\begin{aligned} \frac{58}{7 - 2\sqrt{5}} &= \frac{58}{7 - 2\sqrt{5}} \times \frac{7 + 2\sqrt{5}}{7 + 2\sqrt{5}} \\ &= \frac{58(7 + 2\sqrt{5})}{(7 - 2\sqrt{5})(7 + 2\sqrt{5})} \\ &= \frac{58(7 + 2\sqrt{5})}{(7)^2 - (2\sqrt{5})^2} \\ &= \frac{58(7 + 2\sqrt{5})}{49 - 20} \\ &= \frac{58(7 + 2\sqrt{5})}{29} \\ &= 2(7 + \sqrt{5}) \end{aligned}$$

(ix) Factorize:  $24x^2 - 65x + 21$

**Ans**

$$\begin{aligned} &24x^2 - 65x + 21 \\ &= 24x^2 - 56x - 9x + 21 \\ &= 8x(3x - 7) - 3(3x - 7) \end{aligned}$$



$$= (3x - 7)(8x - 3)$$

3. Write short answers to any Six (6) questions: 12

(i) Find the H.C.F of the following expression:  
 $102xy^2z$ ,  $85x^2yz$  and  $187xyz^2$

**Ans**

$$\begin{aligned} \text{Factors of } 102xy^2z &= 2 \times 3 \times 17 \times x \times y \times y \times z \\ \text{Factors of } 85x^2yz &= 5 \times 17 \times x \times x \times y \times z \\ \text{Factors of } 187xyz^2 &= 11 \times 17 \times x \times y \times z \times z \end{aligned}$$

$$\begin{aligned} \text{H.C.F} &= \text{Multiplication of common factors} \\ &= 17xyz \end{aligned}$$

(ii) Solve the equation:  $\sqrt{5x - 7} - \sqrt{x + 10} = 0$

**Ans**  $\sqrt{5x - 7} - \sqrt{x + 10} = 0$

$$\sqrt{5x - 7} = \sqrt{x + 10}$$

Squaring both sides, we get

$$(\sqrt{5x - 7})^2 = (\sqrt{x + 10})^2$$

$$5x - 7 = x + 10$$

$$5x - x = 10 + 7$$

$$4x = 17$$

$$x = \frac{17}{4}$$

(iii) Solve:  $|2x + 3| = 11$

**Ans**  $|2x + 3| = 11$

$$\pm(2x + 3) = 11$$

$$2x + 3 = 11$$

$$2x = 11 - 3$$

$$2x = 8$$

$$x = \frac{8}{2}$$

$$x = 4$$

$$-(2x + 3) = 11$$

$$2x + 3 = -11$$

$$2x = -11 - 3$$

$$2x = -14$$

$$x = \frac{-14}{2}$$

$$x = -7$$

(iv) Find the value of m and c of  $2x - y = 7$  by expressing it in the form of  $y = mx + c$ .

**Ans** Given,  $2x - y = 7$

$$-y = -2x + 7$$

$$y = 2x - 7$$

(1)



By comparing equ (1) with  $y = mx + c$ , we get

$$m = 2$$

and

$$c = -7$$

(v) Define origin.

**Ans** The point O, where x-axis and y-axis meet is called origin.

(vi) Find the distance between the points:

$$A(-8, 1), B(6, 1)$$

**Ans**  $A(-8, 1), B(6, 1)$

$$\text{Here, } x_1 = -8, y_1 = 1$$

$$x_2 = 6, y_2 = 1$$

The Distance Formula is:

$$\begin{aligned} d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\ &= \sqrt{(6 - (-8))^2 + (1 - 1)^2} \\ &= \sqrt{(6 + 8)^2 + (0)^2} \\ &= \sqrt{(14)^2} \\ &= 14 \end{aligned}$$

(vii) Define scalene triangle.



**Ans** A triangle is called a scalene triangle if measures of all the three sides are different.

(viii) State S.A.S. postulate.

**Ans** In any correspondence of two triangles, if two sides and their included angle of one triangle are congruent to the corresponding two sides and their included angle of the other triangle, then the triangles are congruent.

In  $\triangle ABC \leftrightarrow \triangle DEF$ , shown in the following figures,

$$\text{if } \begin{cases} \overline{AB} \cong \overline{DE} \\ \angle A \cong \angle D \\ \overline{AC} \cong \overline{DF} \end{cases}$$



then  $\triangle ABC \cong \triangle DEF$

(S.A.S Postulate)



Define parallelogram.

**(ix) Ans** A figure formed by four non-collinear points in the plane is called a parallelogram if:

1. its opposite sides are of equal measure;
2. its opposite sides are parallel;
3. measure of none of the angles is  $90^\circ$ .

4. Write short answers to any Six (6) questions: 12

**(i) Define the bisector of a line segment.**

**Ans** A line 'l' is called the bisector of line segment if l is perpendicular to the line segment and passes through its mid-point.

**(ii) 3 cm, 4 cm, 5 cm are the length of the triangle. Give the reason.**

**Ans**


$3 + 4 > 5$	(i)
$3 + 5 > 4$	(ii)
$4 + 5 > 3$	(iii)

From (i), (ii) and (iii) it is proved that the given set can form a triangle. Because, by theorem, the sum of the lengths of any two sides of a triangle is greater than the length of the third side.

**(iii) Define congruent triangles.**

**Ans** Two triangles are said to be congruent written symbolically as,  $\cong$ , if there exists a correspondence between them such that all the corresponding sides and angles are congruent.

**(iv) Find unknown value of x in given figure:**



**Ans** As the above triangle is right angled  $\triangle ABC$ . So, In right angled, by Pythagoras Theorem:

$$(\overline{AC})^2 = (\overline{AB})^2 + (\overline{BC})^2$$

$$(\sqrt{2})^2 = (x)^2 + (1)^2$$

$$2 = x^2 + 1$$




$$\begin{aligned} 2 - 1 &= x^2 \\ \Rightarrow x^2 &= 1 \\ \sqrt{x^2} &= \sqrt{1} \\ x &= 1 \text{ cm} \end{aligned}$$

(v) What is converse of Pythagoras theorem?

**Ans** If the square of one side of a triangle is equal to the sum of squares of the other two sides then the triangle is a right angled triangle.

(vi) Find area of given figure:



**Ans** Length of the rectangle = 6 cm  
 Width of the rectangle = 3 cm  
 Area of the rectangle = Length  $\times$  Width  
 $= 6 \times 3$   
 $= 18 \text{ sq. cm}$

(vii) Define the triangular region.

**Ans** A triangular region is the union of a triangle and its interior, i.e., the three line segments forming the triangle and its interior.


(viii) What is meant by circumcentre?

**Ans** The point of concurrency of the three perpendicular bisectors of the sides of a triangle is called the circumcentre of the triangle.

(ix) Construct a  $\triangle ABC$  in which:

$$m\overline{AB} = 4.2 \text{ cm}, m\overline{BC} = 3.9 \text{ cm}, m\overline{CA} = 3.6 \text{ cm}$$

**Ans**





## Steps of Construction:

1. Take a line segment  $\overline{AB}$  of length 4.2 cm.
2. Take A as centre and draw an arc of 3.6 cm radius.
3. Take B as centre and draw an arc of 3.9 cm radius. This cuts the first arc at C.
4. Join C to A, B.  
ABC is the required triangle.

## (Part-II)

**NOTE:** Attempt THREE (3) questions in all. But question No. 9 is Compulsory.

Q.5.(a) Solve the system of linear equations by Cramer's rule: (4)

$$2x - 2y = 4$$

$$3x + 2y = 6$$

**Ans** For Answer see Paper 2017 (Group-I), Q.5.(a).

(b) Simplify:  $\left(\frac{a^{2l}}{a^{l+m}}\right) \left(\frac{a^{2m}}{a^{m+n}}\right) \left(\frac{a^{2n}}{a^{n+l}}\right)$  (4)

**Ans** For Answer see Paper 2018 (Group-II), Q.5.(b).

Q.6.(a) Use log table to find the value of: (4)

$$0.8176 \times 13.64$$

**Ans**

$$x = 0.8176 \times 13.64$$

$$\log x = \log (0.8176 \times 13.64)$$

$$= \log 0.8176 + \log 13.64$$

$$= 0.0874 + 1.1348$$

$$\log x = 1.0473$$

$$\text{Antilog}(\log x) = \text{Antilog}(1.0473)$$

$$x = 11.15$$

(b) If  $m + n + p = 10$  and  $mn + np + mp = 27$ , then find the value of  $m^2 + n^2 + p^2$ . (4)

**Ans** For Answer see Paper 2017 (Group-I), Q.6.(b).

Q.7.(a) Factorize:  $9x^4 + 36y^4$  (4)

**Ans**  $9x^4 + 36y^4 = 9x^4 + 36y^4 + 36x^2y^2 - 36x^2y^2$   
 $= (3x^2)^2 + (6y^2)^2 + 2(3x^2)(6y^2) - (6xy)^2$   
 $= (3x^2 + 6y^2)^2 - (6xy)^2$



$$= (3x^2 + 6y^2 + 6xy)(3x^2 + 6y^2 - 6xy)$$

$$= (3x^2 + 6xy + 6y^2)(3x^2 - 6xy + 6y^2)$$

- (b) For what value of  $k$  is  $(x + 4)$  the H.C.F of  $x^2 + x - (2k + 2)$  and  $2x^2 + kx - 12$ ? (4)

**Ans** For Answer see Paper 2016 (Group-I), Q.7.(b).

Q.8.(a) Solve:  $-5 \leq \frac{4 - 3x}{2} < 1$  (4)

**Ans** Firstly, multiply by 2

$$-10 \leq 4 - 3x < 2$$

Subtracting by '4' we get

$$-10 - 4 \leq 4 - 3x - 4 < 2 - 4$$

$$-14 \leq -3x < -2$$

Dividing by  $-3$ , we have

$$\frac{-14}{-3} \geq \frac{-3x}{-3} > \frac{-2}{-3} \quad (\text{Change of Sign})$$

$$\frac{14}{3} \geq x > \frac{2}{3}$$

$$\frac{2}{3} < x \leq \frac{14}{3}$$

- (b) Construct the  $\triangle ABC$ , also draw the bisectors of their angles: (4)

$$m\overline{AB} = 3.6 \text{ cm}, m\overline{BC} = 4.2 \text{ cm and } m\angle B = 75^\circ$$

**Ans** For Answer see Paper 2016 (Group-I), Q.8.(b).

- Q.9. Prove that any point inside an angle, equidistant from its arms, is on the bisector of it. (8)

**Ans** For Answer see Paper 2015 (Group-I), Q.9.


OR

Prove that parallelograms on the same base and between the same parallel line (or of the same altitude) are equal in area.

**Ans** Given:

Two parallelograms ABCD and ABEF having the same base  $\overline{AB}$  and between the same parallel lines  $\overline{AB}$  and  $\overline{DE}$ .





**To Prove:**

The area of parallelogram ABCD = area of parallelogram ABEF.

**Proof:**

Statements	Reasons
area of (parallelogram ABCD) = area of (quad. ABED) + area of ( $\Delta$ CBE) (1)	[Area addition axiom]
area of (parallelogram ABEF) = area of (quad. ABED) + area of ( $\Delta$ DAF) (2)	[Area addition axiom]
In $\Delta$ s CBE and DAF $\overline{mCB} = \overline{mDA}$  $\overline{mBE} = \overline{mAF}$  $\angle CBE = \angle DAF$ $\Delta CBE \cong \Delta DAF$	[opposite sides of a parallelogram] [opposite sides of a parallelogram] [ $\because BC \parallel AD, BE \parallel AF$ ]
$\therefore$ area of ( $\Delta$ CBE) = area of ( $\Delta$ DAF) (3) Hence area of (parallelogram ABCD) = area of (parallelogram ABEF)	[cong. area axiom]  from (1), (2) and (3)