

9th Class 2018

Math (Science)

Group-II

Paper-I

Time: 20 Minutes

(Objective Type)

Max Marks: 15

Note: Four possible answers A, B, C and D to each question are given. The choice which you think is correct, fill that circle in front of that question with Marker or Pen ink in the answer-book. Cutting or filling two or more circles will result in zero mark in that question.

1-1- Order of transpose of $\begin{bmatrix} 2 & 1 \\ 0 & 1 \\ 3 & 2 \end{bmatrix}$ is:

- (a) $3 - by - 2 \checkmark$ (b) $1 - by - 2$
 (c) $2 - by - 1$ (d) $2 - by - 3$

2- Every real number is:

- (a) A positive integer
 (b) A rational number
 (c) A negative integer
 (d) A complex number \checkmark

3- $\log e = \text{----}$, where ($e \approx 2.718$):

- (a) 0 (b) $0.4343 \checkmark$
 (c) ∞ (d) 1

4- Conjugate of surd $a + \sqrt{b}$ is ----:

- (a) $-a + \sqrt{b}$ (b) $a - \sqrt{b} \checkmark$
 (c) $\sqrt{a} + \sqrt{b}$ (d) $\sqrt{a} - \sqrt{b}$

5- Find 'm' so that $x^2 + 4x + m$ is a complete square:

- (a) 8 (b) -8
 (c) $4 \checkmark$ (d) 16

6- The square root of $a^2 - 2a + 1$ is ----:

- (a) $\pm(a + 1)$ (b) $\pm(a - 1) \checkmark$
 (c) $a - 1$ (d) $a + 1$

If x is no larger than 10, then ---- :

- 7- (a) $x \geq 8$ (b) $x \leq 10$ ✓
(c) $x < 10$ (d) $x > 10$
- 8- Point (2, -3) lies in quadrant:
(a) I (b) II
(c) III (d) IV ✓
- 9- Mid-point of the points (0, 0) and (2, 2) is:
(a) (1, 1) ✓ (b) (1, 0)
(c) (0, 1) (d) (-1, -1)
- 10- Notation used for congruent is:
(a) \perp (b) $=$
(c) \cong ✓ (d) \sim
- 11- Notation \parallel means:
(a) Congruent (b) Equal
(c) Parallel ✓ (d) Un-equal
- 12- Bisection means to divide into equal parts:
(a) 1 (b) 3
(c) 4 (d) 2 ✓
- 13- The unit of ratio is:
(a) kg (b) m
(c) cm (d) None of these ✓
- 14- A ray has ---- end points:
(a) 2 (b) 1 ✓
(c) 3 (d) 4
- 15- The right bisectors of the three sides of a triangle are ---- :
(a) Congruent (b) Collinear
(c) Concurrent ✓ (d) Parallel

9th Class 2018

Math (Science)	Group-II	Paper-I
Time: 2.10 Hours	(Subjective Type)	Max. Marks: 60

(Part-I)

2. Write short answers to any Six (6) questions: 12

(i) Define transpose of matrix.

Ans A matrix obtained by changing the row into columns or columns into rows of a matrix is called transpose of that matrix. If A is a matrix, then its transpose is denoted by A^t .

(ii) Find additive inverse of the matrices:

$$\begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$$

Ans Let:

$$A = \begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$$

Then additive inverse of A is:

$$A = \begin{bmatrix} -\sqrt{3} & -1 \\ 1 & -\sqrt{2} \end{bmatrix}$$

(iii) Define multiplicative identity.

Ans Let A be a matrix. Another matrix B is called the identity matrix of A under multiplication if

$$AB = A = BA$$

(iv) Simplify: $5^{2^3} \div (5^2)^3$

Ans

$$= 5^{2^3} \div (5^2)^3$$

$$= 5^8 \div 5^6$$

$$= 5^{8-6}$$

$$= 5^2$$

$$\boxed{= 25}$$

(v) Find the value of x , when: $\log_{64} 8 = \frac{x}{2}$

Ans

$$\log_{64} 8 = \frac{x}{2}$$

$$(64)^{x/2} = 8$$

$$(8^2)^{x/2} = 8^1$$

$$8^x = 8^1$$

$$\boxed{x = 1}$$

(vi) Define logarithm.

Ans If $a^x = y$, then x is called the logarithm of y to the base ' a ' and is written as $\log_a y = x$, where $a > 0$, $a \neq 1$ and $y > 0$.

(vii) Simplify: $\frac{x+2}{2x-3y} \cdot \frac{4x^2-9y^2}{xy+2y}$

Ans

$$\begin{aligned} \frac{x+2}{2x-3y} \cdot \frac{4x^2-9y^2}{xy+2y} &= \frac{(x+2)[(2x)^2-(3y)^2]}{(2x-3y)(x+2)y} \\ &= \frac{\cancel{(x+2)}(2x+3y)\cancel{(2x-3y)}}{\cancel{(2x-3y)}\cancel{(x+2)}y} \end{aligned}$$

$$\boxed{= \frac{2x+3y}{y}}$$

(viii) Rationalize the denominator of $\frac{1}{3+2\sqrt{5}}$.

Ans

$$= \frac{1}{3+2\sqrt{5}} \times \left(\frac{3-2\sqrt{5}}{3-2\sqrt{5}} \right)$$

$$= \frac{3-2\sqrt{5}}{(3)^2-(2\sqrt{5})^2}$$

$$= \frac{3-2\sqrt{5}}{9-20}$$

$$= \frac{3-2\sqrt{5}}{-11}$$

$$\boxed{= \frac{-1}{11} (3-2\sqrt{5})}$$

(ix) What is meant by remainder theorem?

Ans If a polynomial $p(x)$ is divided by a linear divisor $(x - a)$, then the remainder is $p(a)$.

3. Write short answers to any Six (6) questions: 12

(i) Find H.C.F. of the polynomials by factorization:

$$x^2 + 5x + 6, x^2 - 4x - 12$$

Ans $x^2 + 5x + 6 = x^2 + 2x + 3x + 6$
 $= x(x + 2) + 3(x + 2) \Rightarrow (x + 2)(x + 3)$

$$x^2 - 4x - 12 = x^2 - 6x + 2x - 12$$

$$= x(x - 6) + 2(x - 6) \Rightarrow (x - 6)(x + 2)$$

$$\text{H.C.F} = x + 2 \text{ (common factor)}$$

(ii) Solve the equation: $\sqrt{3x + 4} = 2$

Ans $(\sqrt{3x + 4})^2 = (2)^2$
 $3x + 4 = 4$

$$3x = 4 - 4$$

$$3x = 0$$

$$\boxed{x = 0}$$

(iii) Find the solution set of: $|3x - 5| = 4$

Ans

$$3x - 5 = 4; \quad 3x - 5 = -4$$

$$3x = 4 + 5; \quad 3x = -4 + 5$$

$$3x = 9; \quad 3x = 1$$

$$x = \frac{9}{3}; \quad x = \frac{1}{3}$$

$$x = 3$$

(iv) Define Cartesian plane.

Ans The Cartesian plane establishes one-to-one correspondence between the set of ordered pairs $R \times R = \{(x, y) \mid x, y \in R\}$ and the points of the Cartesian plane.

(v) Find the value of m and c of the line expressing it in the form $y = mx + c$, $3 - 2x + y = 0$.

Ans

$$y = mx + c \quad (i)$$

$$3 - 2x + y = 0$$

$$y = 2x - 3 \quad (ii)$$

By comparing both equations, we get

$$m = 2,$$

$$c = -3$$

- (vi) Find the distance between pair of points:
A(0, 0), B(0, -5)

Ans

$$\begin{aligned} d &= \sqrt{|x_2 - x_1|^2 + |y_2 - y_1|^2} \\ |AB| &= \sqrt{[(0 - 0)]^2 + [(-5) - 0]^2} \\ &= \sqrt{0 + (-5)^2} \\ &= \sqrt{25} \\ &= 5 \end{aligned}$$

- (vii) Find the mid-point between the pair of points:
A(-4, 9), B(-4, -3)

Ans


$$\begin{aligned} &A(-4, 9), B(-4, -3) \\ P(x, y) &= \left(\frac{-4 - 4}{2}, \frac{9 - 3}{2} \right) \\ P(x, y) &= (-4, 3) \\ \text{Mid-point of AB} &= (-4, 3) \end{aligned}$$

- (viii) What is meant by the congruency of triangles?

Ans Two triangles are said to be congruent, if there exists a correspondence between them such that all the corresponding sides and angles are congruent.

- (ix) One angle of a parallelogram is 130° . Find the measures of its remaining angles.

Ans



$$\begin{aligned} \angle B &\cong \angle C \\ m\angle A &= 130^\circ \\ m\angle C &= 130^\circ \\ m\angle B &= 180^\circ - m\angle A \\ &= 180^\circ - 130^\circ = 50^\circ \end{aligned}$$

As

$$\begin{aligned} \angle B &= \angle D \\ m\angle C &= 50^\circ \end{aligned}$$

4. Write short answers to any Six (6) questions: 12
- (i) If 3 cm and 4 cm are lengths of two sides of a right angle triangle, then what should be the third length of the triangle?

Ans

$$(\text{Hypotenuse})^2 = (\text{Perpendicular})^2 + (\text{Base})^2$$

$$(\text{Hypotenuse})^2 = (3)^2 + (4)^2$$

$$(\text{Hypotenuse})^2 = 9 + 16$$

$$(\text{Hypotenuse})^2 = 25$$

$$\sqrt{(\text{Hypotenuse})^2} = \sqrt{25}$$

$$(\text{Hypotenuse}) = 5 \text{ cm}$$

- (ii) Define bisector of an angle.

Ans

Angle bisector is the ray which divides an angle into two equal parts.

- (iii) Define proportion.

Ans

Equality of two ratios is defined as the proportion.


if $a : b = c : d$, then a, b, c and d are said to be a proportion.

- (iv) State converse to Pythagoras theorem.

Ans

Converse of Pythagoras theorem is:

If the square of one side of a triangle is equal to the sum of the squares of the other two sides, then the triangle is a right-angled triangle.



- (v) Verify that the triangle having the measures of sides $a = 1.5 \text{ cm}$, $b = 2 \text{ cm}$, $c = 2.5 \text{ cm}$ are right-angled.

Ans

$$a = 1.5 \text{ cm}, b = 2 \text{ cm}, c = 2.5 \text{ cm}$$

$$c^2 = a^2 + b^2$$

$$(2.5)^2 = (1.5)^2 + (2)^2$$

$$6.25 = 2.25 + 4$$


$$6.25 = 6.25$$

Hence measures are the sides of a triangle.

(vi) Define rectangular region.

Ans A rectangular region is the union of a rectangle and its interior.

(vii) Find the area of the figure:



Ans Length of rectangle = 6 cm

Width of // // = 3 cm

Area of // // = 6×3

= 18 Sq. cm


(viii) Define incentre.

Ans The internal bisectors of the angles of a triangle meet at a point called the incentre of the triangle.

(ix) Construct a $\triangle ABC$ in which:

$m\overline{AB} = 3.2 \text{ cm}$, $m\overline{BC} = 4.2 \text{ cm}$, $m\overline{CA} = 5.2 \text{ cm}$

Ans



(Part-II)

NOTE: Attempt THREE questions in all. But question No. 9 is Compulsory.

Q.5.(a) Solve with the help of Cramer's rule:

(4)

$$2x + y = 3$$

$$6x + 5y = 1$$

Ans

$$2x + y = 3$$

$$6x + 5y = 1$$

$$A = \begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix}$$

$$Ax = \begin{bmatrix} 3 & 1 \\ 1 & 5 \end{bmatrix}$$

$$Ay = \begin{bmatrix} 2 & 3 \\ 6 & 1 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & 1 \\ 6 & 5 \end{vmatrix}$$

$$= (2)(5) - (1)(6)$$

$$= 10 - 6$$

$$= 4$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 3 & 1 \\ 1 & 5 \end{vmatrix}}{4}$$

$$= \frac{(3)(5) - (1)(1)}{4}$$

$$= \frac{15 - 1}{4}$$

$$= \frac{14}{4}$$

$$x = \frac{7}{2}$$

$$y = \frac{|A_y|}{|A|}$$

$$y = \frac{\begin{vmatrix} 2 & 3 \\ 6 & 1 \end{vmatrix}}{4}$$

$$y = \frac{(2)(1) - (6)(3)}{4}$$

$$y = \frac{2 - 18}{4}$$

$$y = \frac{-16}{4}$$

$$y = -4$$

$$x = \frac{7}{2}, y = -4$$

(b) Simplify: $\left(\frac{a^{2l}}{a^{l+m}}\right)\left(\frac{a^{2m}}{a^{m+n}}\right)\left(\frac{a^{2n}}{a^{n+l}}\right)$ (4)

Ans

$$\begin{aligned} &= \left(\frac{a^{2l}}{a^{l+m}}\right)\left(\frac{a^{2m}}{a^{m+n}}\right)\left(\frac{a^{2n}}{a^{n+l}}\right) \\ &= a^{2l-(l+m)} \times a^{2m-(m+n)} \times a^{2n-(n+l)} \\ &= a^{2l-l-m} \times a^{2m-m-n} \times a^{2n-n-l} \\ &= a^{l-m} \times a^{m-n} \times a^{n-l} \\ &= a^{l-m+m-n+n-l} \\ &= a^0 \end{aligned}$$

$$\boxed{= 1}$$

Q.6.(a) Use log table to find the value of : (4)

$$\frac{0.678 \times 9.01}{0.0234}$$

Ans Let,

$$x = \frac{0.678 \times 9.01}{0.0234}$$

Taking log both side

$$\begin{aligned} \log x &= \log \frac{0.678 \times 9.01}{0.0234} \\ &= \log 0.678 + \log 9.01 - \log 0.0234 \\ &= \bar{1}.8312 + 0.9547 - (\bar{2}.3692) \\ &= \bar{1}.8312 + 0.9547 - \bar{2}.3692 \\ &= -1 + .8312 + 0.9547 + 2 - .3692 \\ &= 2.4167 \end{aligned}$$

Take antilog

$$x = \text{Antilog } 2.4167$$

$$\boxed{x = 261}$$

(b) If $x + y = 7$ and $xy = 12$, then find the value of $x^3 + y^3$. (4)

Ans

$$\begin{aligned} x + y &= 7 \\ xy &= 12 \\ x^3 + y^3 &= ? \end{aligned}$$

Formula:

$$(x + y)^3 = x^3 + y^3 + 3xy(x + y)$$

Putting values,

$$(7)^3 = x^3 + y^3 + 3(12)(7)$$

$$343 = x^3 + y^3 + 252$$

$$343 - 252 = x^3 + y^3$$

$$91 = x^3 + y^3$$

$$\boxed{x^3 + y^3 = 91}$$

Q.7.(a) For what value of m is the polynomial

$$p(x) = 4x^3 - 7x^2 + 6x - 3m$$

exactly divisible by $x + 2$?

(4)

Ans

$$P(x) = 4x^3 - 7x^2 + 6x - 3m$$

From $x + 2 = 0$, $x = -2$

$$P(-2) = 4(-2)^3 - 7(-2)^2 + 6(-2) - 3m$$

$$= -32 - 28 - 12 - 3m$$

$$= -72 - 3m$$

If $x + 2$ is factor, then $R = 0$.

$$-72 - 3m = 0$$

$$-3(24 + m) = 0$$

$$24 + m = 0$$

$$\boxed{m = -24}$$

(b) Simplify to the lowest form:

(4)

$$\frac{2y^2 + 7y - 4}{3y^2 - 13y + 4} \div \frac{4y^2 - 1}{6y^2 + y - 1}$$

Ans

$$= \frac{2y^2 + 7y - 4}{3y^2 - 13y + 4} \div \frac{4y^2 - 1}{6y^2 + y - 1}$$

$$= \frac{2y^2 + 8y - y - 4}{3y^2 - 12y - y + 4} \div \frac{(2y)^2 - (1)^2}{6y^2 + 3y - 2y - 1}$$

$$= \frac{2y(y + 4) - 1(y + 4)}{3y(y - 4) - 1(y - 4)} \div \frac{(2y + 1)(2y - 1)}{3y(2y + 1) - 1(2y + 1)}$$

$$= \frac{(2y - 1)(y + 4)}{(3y - 1)(y - 4)} \div \frac{(2y + 1)(2y - 1)}{(3y - 1)(2y + 1)}$$

$$= \frac{(2y - 1)(y + 4)}{(3y - 1)(y - 4)} \times \frac{(3y - 1)}{(2y - 1)}$$

$$= \frac{y+4}{y-4}$$

Q.8.(a) Find the solution set of the equation:

(4)

$$\frac{x}{3x-6} = 2 - \frac{2x}{x-2}, x \neq 2$$

Ans

$$\frac{x}{3x-6} + \frac{2x}{x-2} = 2$$

$$\frac{x}{3(x-2)} + \frac{2x}{x-2} = 2$$

$$\frac{x + 3(2x)}{3(x-2)} = 2$$

$$\frac{x + 6x}{3x-6} = 2$$

$$\frac{7x}{3x-6} = 2$$

$$7x = 2(3x-6)$$

$$7x = 6x - 12$$

$$7x - 6x = -12$$


$$x = \{-12\}$$

(b) Construct $\triangle ABC$. Draw perpendicular bisectors of its sides:

(4)

$$m\angle A = 120^\circ, \quad m\overline{AC} = 3.2 \text{ cm}, \quad m\overline{AB} = 2.4 \text{ cm}$$

Ans



Step of Construction:

- (i) Take $m\overline{AB} = 2.4 \text{ cm}$.
- (ii) Draw $m\angle BAC = 120^\circ$ at point A.

- (iii) With centre at the point A and radius 3.2 cut $\widehat{mAC} = 3.2$ cm.
- (iv) Join B to C to complete the $\triangle ABC$.
- (v) Draw perpendicular bisectors of BC and CA meeting at point O.
- (vi) Now draw perpendicular bisector of third side AB.
- (vii) We observe that it also passes through O, the point of intersection of first two perpendicular bisectors.
- (viii) Hence the three perpendicular bisectors of $\triangle ABC$ are concurrent at O.

Q.9. Prove that the right bisectors of the sides of a triangle are concurrent. (8)

Ans For Answer see Paper 2017 (Group-I), Q.9.

OR

Prove that triangles on equal bases and of equal altitudes are equal in area.

Ans For Answer see 2014 (Group-II), Q.9(OR).